

Equivalence of two thermostatistical formalisms based on the Havrda & Charvat-Daróczy-Tsallis entropies

G.A. Raggio*

Facultad de Matemática, Astronomía y Física
 Universidad Nacional de Córdoba
 Ciudad Universitaria, 5000 Córdoba, Argentina
 (August 14, 1999)

We show that the latest thermostatistical formalism based on the Havrda & Charvat-Daróczy-Tsallis entropy $S_q[\rho] = (q-1)^{-1}(1 - \text{tr}(\rho^q))$ proposed by Tsallis, Mendes and Plastino is *equivalent* to the first one proposed by Tsallis in 1988. Here, equivalent means: *the “equilibrium” state predicted by either formalism using q leads to the same expectation values for all observables as that predicted by the other formalism using $1/q$ ”*. We also point out once again that the basic property of *transitivity of equilibrium* (e.g., the 0th Law of Thermodynamics) fails in these formalisms.

PACS Numbers: 05.30.-d, 05.90.+m

In a recent paper [1], Tsallis, Mendes and Plastino introduce a further thermostatistical formalism based on the Havrda & Charvat-Daróczy-Tsallis entropy [2–4]

$$S_q[\rho] = (q-1)^{-1}(1 - \text{tr}(\rho^q)) \quad , \quad 0 < q \neq 1 . \quad (1)$$

This third version of Tsallis’ thermostatistical formalism proposes to maximize $S_q[\rho]$ among the states ρ for which

$$\langle A \rangle_\rho^{(3)} := \frac{\text{tr}(A\rho^q)}{\text{tr}(\rho^q)} \quad (2)$$

is constant for some (operator) observable A . The purpose of this note is to show that this “new” formalism is something of a red herring: it is completely equivalent to maximizing $S_{1/q}[\rho]$ among the states ρ for which the usual expectation value

$$\langle A \rangle_\rho = \text{tr}(A\rho) \quad (3)$$

is constant.

This last formalism – indicated by the superindex (1) in [1] – is Tsallis’ original proposal of [4]; although in that paper Tsallis misidentifies the “temperature” in the sense that his parameter t is not directly the correct “temperature” arising from the Legendre transformation procedure¹. There is a further formalism – chronologically the second – due to Curado and Tsallis [6] (the superindex (2) is used for this in [1]) which uses the “expectation value” formula

$$\langle A \rangle_\rho^{(2)} = \text{tr}(A\rho^q) \quad (4)$$

which has, among many other drawbacks², the feature that the “expectation value” of a sum is not the sum of the “expectations” and the “expectation” of a constant is not constant (and consequences thereof). This is remedied in [1] by renormalizing (4) to get (2). However, both these “expectations” fail to be affine in the state; i.e., the analogues of the relation

$$\langle A \rangle_{\lambda\rho_1 + (1-\lambda)\rho_2} = \lambda \langle A \rangle_{\rho_1} + (1-\lambda) \langle A \rangle_{\rho_2} \quad (5)$$

for the mixture of the states ρ_1 and ρ_2 with $0 \leq \lambda \leq 1$, fail. This relation is crucial for a consistent probabilistic interpretation.

We now come to our claim. First of all we are using the term “state” to denote a positive trace class operator of unit trace (i.e., a density operator), and “tr” denotes the trace. We choose to discuss these matters in the domain of

*Investigador CONICET; e-mail: raggio@fis.uncor.edu

¹see [5], for a correct introduction of the “temperature” in this formalism along with a detailed description of some of its problems, notably the failure of the O^{th} -law of Thermodynamics.

²These are expounded in [7], and include the failure of the 0th-law of Thermodynamics.

quantum mechanics; the discussion in the domain of classical mechanics is identical (replace “state” by “probability density” and the trace by an appropriate integral or sum). We fix some value of q greater than 0 and not equal to 1.

Associate to each state ρ the state $\mathcal{R}_q(\rho)$ given by

$$\mathcal{R}_q(\rho) := (tr(\rho^q))^{-1} \rho^q ; \quad (6)$$

which has $\mathcal{R}_{1/q}$ as its inverse:

$$\mathcal{R}_q(\mathcal{R}_{1/q}(\rho)) = \mathcal{R}_{1/q}(\mathcal{R}_q(\rho)) = \rho ; \quad (7)$$

and clearly satisfies

$$\langle A \rangle_\rho^{(3)} = \langle A \rangle_{\mathcal{R}_q(\rho)} . \quad (8)$$

When $0 < q < 1$ and in the case of an infinite dimensional underlying Hilbert space some care has to be exercised because $tr(\rho^q)$ and thus $S_q[\rho]$ can be infinite. However, as shown in [8], the set of states with finite q -entropy is convex (and $S_q[\cdot]$ is strictly concave for these states). In what follows we will not go into details of this and simply assume (with no loss of content or generality) that we are dealing with states of finite q -entropy. The following simple observation is the basis of our claim.

Lemma: Let ρ and ω be arbitrary states, then for every $0 < q \neq 1$, one has $S_q[\rho] \leq S_q[\omega]$ if and only if $S_{1/q}[\mathcal{R}_q(\rho)] \leq S_{1/q}[\mathcal{R}_q(\omega)]$.

Proof: A simple algebraic manipulation shows that for arbitrary states ρ and ω one has: $S_q[\rho] \geq S_q[\omega]$ if and only if $tr(\rho^q) \leq tr(\omega^q)$, when $q > 1$; and $S_q[\rho] \geq S_q[\omega]$ if and only if $tr(\rho^q) \geq tr(\omega^q)$, when $q < 1$. Notice also that

$$tr(\mathcal{R}_q(\rho)^{1/q}) = (tr(\rho^q))^{-1/q} .$$

Let ρ and ω be arbitrary states. The inequality $tr(\rho^q) \leq tr(\omega^q)$ is equivalent to the inequality $(tr(\rho^q))^{-1/q} \geq (tr(\omega^q))^{-1/q}$, which is equivalent to the inequality $tr((\mathcal{R}_q(\rho))^{1/q}) \geq tr((\mathcal{R}_q(\omega))^{1/q})$. Using the observation of the previous paragraph to shift this inequality to the entropies, one obtains the claim.

The Tsallis-Mendes-Plastino proposal ([1]) reads: Maximize $S_q[\rho]$ among the states ρ for which $\langle A \rangle_\rho^{(3)} = \alpha$ to find a maximizing state $\varphi_{\alpha,q}$.

My proposal would be: Maximize $S_{1/q}[\rho]$ among the state ρ for which $\langle A \rangle_\rho = \alpha$ to find a unique ³ maximizing state $\mu_{\alpha,1/q}$.

Then eqs. (7) and (8) together with the Lemma say that $\mu_{\alpha,1/q} = \mathcal{R}_q(\varphi_{\alpha,q})$. Moreover, both proposals have the same expectation values, the computation of which never needs $\varphi_{\alpha,q}$ since these are calculated in the usual manner tracing with $\mu_{\alpha,1/q}$. Thus, if for whatever reason, you should like to consider $S_q[\cdot]$ as the underlying entropy then there is no need to use the strange expectation (2) but can proceed with the usual one – albeit replacing q by $1/q$.

Explicit expressions for $\nu_{\alpha,q}$ as well as for the “thermodynamic functions” associated with the map $\alpha \mapsto S_q[\mu_{\alpha,q}]$ via Legendre transformation are given in [5].

I allow myself some comments prompted by reading [1] as the last of many papers on the subject. I find it remarkable that after a considerable amount of publications on the various (mainly the second) Tsallis formalisms (see the up-to-date list in the URL <http://tsallis.cat.cbpf.br/biblio.htm>) there is not a single piece of evidence supporting the claim that these formalisms constitute a generalization of Boltzmann-Gibbs statistical mechanics. It is true that the formalism of statistical mechanics can be obtained by the statistical inference route (championed by Szilard and Jaynes) by maximizing the usual entropy with the usual constraint and identifying the various variables with the corresponding thermodynamic ones. However, Boltzmann’s programme (compare [9]) provides a way of obtaining the results of this variational exercise from the laws of physics. This link is as yet missing for the Tsallis formalisms. I also see severe problems connected with the interpretation of various unusual features of the formalisms based on

³The uniqueness, given in [5], is simple to prove: for given α , if ρ_1 and ρ_2 are distinct maximizers and $S_q[\rho_1] = S_q[\rho_2] = s < \infty$, then since (3) is an affine constraint (5), the state $\lambda\rho_1 + (1-\lambda)\rho_2$ has $\langle A \rangle_{\lambda\rho_1 + (1-\lambda)\rho_2} = \alpha$ for any $0 < \lambda < 1$. Since $S_q[\cdot]$ is strictly concave, one obtains a contradiction via $S_q[\lambda\rho_1 + (1-\lambda)\rho_2] > \lambda S_q[\rho_1] + (1-\lambda)S_q[\rho_2] = s$.

S_q , notably the failure of the 0^{th} Law or more generally, the fact that the intensive parameters derived by Legendre transformation are not “equilibrium” parameters in the sense that they are not “tags” for a *transitive* equilibrium relation. This feature of the formalisms was pointed out in [5,7].

Let me present once again the results of [5] (where the proofs are detailed) and the argument leading to this disturbing feature. For fixed A (say the energy observable) consider the entropy function

$$S_q(\alpha) = \sup_{\{\rho: \langle A \rangle_\rho = \alpha\}} S_q[\rho] . \quad (9)$$

along with its Legendre-Fenchel transform

$$\phi_q(\beta) = \inf_{\alpha} \{\beta\alpha - S_q(\alpha)\} = \inf_{\rho} \{\beta \langle A \rangle_\rho - S_q[\rho]\} . \quad (10)$$

Then $S_q(\cdot)$ is the Legendre-Fenchel transform of $\phi_q(\cdot)$ – a fact that really closes the circle leading to a full-fledged “thermal” formalism – and there is a unique state⁴ $\mu_{\alpha,q}$ such that $S_q[\mu_{\alpha,q}] = S_q(\alpha)$; there is a unique state $\nu_{\beta,q}$ minimizing the problem (10); one has $\mu_{\alpha,q} = \nu_{\beta(\alpha),q}$ where the function $\alpha \mapsto \beta(\alpha)$ is determined by the relation $(d\phi_q/d\beta)(\beta(\alpha)) = \alpha$.

The formulas for the state $\nu_{\beta,q}$ are given explicitly in [5] for the various cases: finite/infinite dimension, and $0 < q < 1/q > 1$.

In the most interesting case of infinite dimension with A bounded below with purely discrete spectrum $\{a_n : n = 0, 1, 2, \dots\}$, numbered in order of increasing order not counting the multiplicities m_n (assumed finite), and such that

$$\sum_{n \geq 1} m_n (a_n - a_o)^{q/(q-1)} < \infty ,$$

one has

$$\nu_{\beta,q} = \frac{(1 + (1-q)\tau_q(\beta)(A - a_o))^{1/(q-1)}}{\text{trace of the above}} \quad (11)$$

for all positive⁵ β when $0 < q < 1$.

For $q > 1$ and always in the infinite dimensional case with A bounded below with purely discrete spectrum but no summability condition, one has

$$\nu_{\beta,q} = \frac{[(1 + (1-q)\tau_q(\beta)(A - a_o))]_{\oplus}^{1/(q-1)}}{\text{trace of the above}} \quad (12)$$

for all β in a certain explicitly given finite subinterval $(0, \beta_c)$ of the positive reals⁶. Above β_c the state $\nu_{\beta,q}$ is the equidistribution of ground states and independent of β ⁷. In the last formula, $[\dots]_{\oplus}$ means “take the positive part of the operator ...”. In both formulas $\tau_q(\beta)$ is a certain implicitly determined, strictly increasing function depending on A (see [5]). Notice that $\nu_{\beta,q}$ is always degenerate since all eigenstates of A above a certain energy depending on β are not populated!

If one considers two subsystems 1, and 2 characterized by their respective Hamiltonians A_1 and A_2 , of one quantum system with the Hamiltonian $A = A_1 \otimes \mathbf{1} + \mathbf{1} \otimes A_2 + \lambda V$. Boltzmann-Gibbs statistical mechanics predicts

$$\frac{\exp(-\beta A)}{\text{tr}(\exp(-\beta A))}$$

as the equilibrium state to the intensive parameter β . One has

$$\frac{\exp(-\beta A)}{\text{tr}(\exp(-\beta A))} \rightarrow \frac{\exp(-\beta A_1)}{\text{tr}_1(\exp(-\beta A_1))} \otimes \frac{\exp(-\beta A_2)}{\text{tr}_2(\exp(-\beta A_2))}$$

⁴In the infinite dimensional situation and for $0 < q < 1$, restrictions on the spectrum of A have to be imposed in order to get finite solutions.

⁵ $\phi_q(\beta) = -\infty$ for $\beta \leq 0$ so that negative “temperatures” do not arise.

⁶ $\phi_q(\beta) = -\infty$ for $\beta \leq 0$.

⁷There is thus a minimal possible “temperature”.

in the limit of weak coupling $\lambda \rightarrow 0$. This is the product-state formed with the equilibrium states of the subsystems for the same value of β . This is the fact needed to obtain transitivity of β -equilibrium: two systems in β -equilibrium with a third are in β -equilibrium with each other.

The reader is asked to insert $A = A_1 \otimes \mathbf{1} + \mathbf{1} \otimes A_2 + \lambda V$ in either of the formulas (11) or (12) and to see that when $\lambda \rightarrow 0$ one does not obtain a product state.

I am indebted to Pedro Pury for drawing my attention to [1], and to him and Gustavo Guerberoﬀ with whom most of my work on the S_q -formalisms was shared in 1995. The support of CONICET, CONICOR and SECYT-UNC is gratefully acknowledged.

-
- [1] C. Tsallis, R.S. Mendes and A.R. Plastino: *The role of constraints within generalized nonextensive statistics*. Physica A **261**, 534-554 (1998).
 - [2] J. Havrda and F. Charvat: *Quantification method of classification process: concept of structural α -entropy*, Kybernetika 3, 30 (1967). I am indebted to Guy Jumaire for pointing this reference out to me in August 1996.
 - [3] Z. Daróczy: *Generalized information functions*. Inform. Control **16**, 36-51 (1970).
 - [4] C. Tsallis: *Possible generalization of Boltzmann-Gibbs Statistics*. J. Stat. Phys. **52**, 479-487 (1988).
 - [5] G. Guerberoﬀ, and G.A. Raggio: *Standard thermal statistics with q -entropies*. J. Math. Phys. **37**, 1776-1789 (1996).
 - [6] E.M.F. Curado, and C. Tsallis: *Generalized statistical mechanics: connection with thermodynamics*. J. Phys. A **24**, L69-L72 (1991).
 - [7] G. Guerberoﬀ, P.A. Pury, and G.A. Raggio: *Nonstandard thermal statistics with q -entropies*. J. Math. Phys. **37**, 1790-1811 (1996).
 - [8] G.A. Raggio: *Properties of q -entropies*. J. Math. Phys. **36**, 4785-4791 (1995).
 - [9] J.L. Lebowitz: *Boltzmann's Entropy and Time's Arrow*. Physics Today **46** (9), 32-38 (1993).